



# VERTEX EQUITABLE LABELING OF UNION OF CYCLIC SNAKE IN TRANSFORMED TREES

A. Maheswari<sup>1</sup> , K.Thanalakshmi<sup>2</sup>

<sup>1,2</sup> Department of Mathematics, Kamaraj College of Engineering and Technology,  
Virudhunagar, (India)

## ABSTRACT

Let  $G$  be a graph with  $p$  vertices and  $q$  edges and  $A = \left\{0, 1, 2, \dots, \left\lceil \frac{q}{2} \right\rceil\right\}$ . A vertex labeling  $f: V(G) \rightarrow A$  induces an edge labeling  $f^*$  defined by  $f^*(uv) = f(u) + f(v)$  for all edges  $uv$ . For  $a \in A$ , let  $v_f(a)$  be the number of vertices  $v$  with  $f(v) = a$ . A graph  $G$  is said to be vertex equitable if there exists a vertex labeling  $f$  such that for all  $a$  and  $b$  in  $A$ ,  $\left|v_f(a) - v_f(b)\right| \leq 1$  and the induced edge labels are  $1, 2, 3, \dots, q$ . In this paper, we prove that  $T \hat{\circ} nC_4$  and zig-zag triangle are vertex equitable graphs.

**Key Words:** Vertex Equitable Labeling, Vertex Equitable Graph

**AMS Classification (2010):** 05C78

## I INTRODUCTION

All graphs considered here are simple, finite, connected and undirected. We follow the basic notations and terminologies of graph theory as in [1]. A graph labeling is an assignment of integers to the vertices or edges or both, subject to certain conditions. There are several types of labeling and a detailed survey of graph labeling can be found in [2]. The vertex set and the edge set of a graph are denoted by  $V(G)$  and  $E(G)$  respectively. The concept of vertex equitable labeling was due to Lourdasamy and Seenivasan in [3] and further studied in [4-12]. Let  $G$  be a


graph with  $p$  vertices and  $q$  edges and  $A = \left\{0, 1, 2, \dots, \left\lceil \frac{q}{2} \right\rceil\right\}$ . A graph  $G$  is said to be vertex equitable if there exists a

vertex labeling  $f: V(G) \rightarrow A$  that induces an edge labeling  $f^*$  defined by  $f^*(uv) = f(u) + f(v)$  for all edges  $uv$  such

We use the following definitions in the subsequent section.

**Definition 1.3. [13]** A  $nC_k$ -snake is defined as a connected graph in which all the  $n$ -blocks are isomorphic to the cycle  $C_k$  and the block-cut point graph is a path. Let  $P$  be the path of minimum length that contains all the cut vertices of a  $nC_k$ -snake. Any  $nC_k$ -snake can be represented by a string  $s_1, s_2, \dots, s_{n-2}$  of integers of length  $n-2$  where the  $i^{\text{th}}$  integer,  $s_i$ , on the string is the distance between  $i^{\text{th}}$  and  $(i+1)^{\text{th}}$  cut vertices on the path  $P$  from one extreme and is taken from  $S_k = \{1, 2, \dots, \left\lfloor \frac{k}{2} \right\rfloor\}$ . The strings obtained for both extremes are assumed to be the same. For example,


the string of a  $10C_4$ -snake is shown in Figure 1.1 is 2,2,1,2,1,1,2,1.



A  $nC_k$ –snake is said to be linear if each integer of its string is  $\left\lfloor \frac{k}{2} \right\rfloor$ . The  $nC_4$ –snake graph with diagonal vertices are  $u_{1j}$  ( $1 \leq j \leq n+1$ ), left to the diagonal vertices are  $v_{1j}$  ( $1 \leq j \leq n$ ) and right to the diagonal vertices are  $w_{1j}$  ( $1 \leq j \leq n$ ).

**Definition 1.4:** Let  $H$  be any graph with  $n$  vertices  $v_1, v_2, \dots, v_n$  and let  $G_1, G_2, \dots, G_n$  be  $n$  graphs. Then  $H \hat{O} (G_1, G_2, \dots, G_n)$  is a graph obtained by identifying a vertex  $u_i$  of  $G_i$  with a vertex  $v_i$  of  $H$  for  $1 \leq i \leq n$ . If all the graphs  $G_i$  are isomorphic to a graph  $G$ , then the graph is denoted by  $H \hat{O} G$ . Again  $H \tilde{O} (G_1, G_2, \dots, G_n)$  is a graph obtained by joining a vertex  $u_i$  of  $G_i$  with a vertex  $v_i$  of  $H$  by an edge for  $1 \leq i \leq n$ . If all the graphs  $G_i$  are isomorphic to a graph  $G$ , then the graph is denoted by  $H \tilde{O} G$ .

**Definition 1.3.** Let  $G$  be a graph obtained from the path  $P_n; x_1 x_2 \dots x_n$  adding a new vertices  $y_1 y_2 \dots y_n$  and new edges  $y_1 x_2, y_n x_{n-1}; x_i y_i$  for  $1 \leq i \leq n$ .  $v \cdot x_{i-1}, y_i x_{i+1}$  for  $2 \leq i \leq n-1$ . The family of graphs is called zig-zag triangle.



**Definition 1.4[14].** Let  $T$  be a tree and  $u_0$  and  $v_0$  be the two adjacent vertices in  $T$ . Let  $u$  and  $v$  be the two pendant vertices of  $T$  such that the length of the path  $u_0-u$  is equal to the length of the path  $v_0-v$ . If the edge  $u_0 v_0$  is deleted from  $T$  and  $u$  and  $v$  are joined by an edge  $uv$ , then such a transformation of  $T$  is called an elementary parallel transformation (or an ept) and the edge  $u_0 v_0$  is called transformable edge.

If by the sequence of ept's,  $T$  can be reduced to a path, then  $T$  is called a  $T_p$ -tree (transformed tree) and such sequence regarded as a composition of mappings (ept's) denoted by  $P$ , is called a parallel transformation of  $T$ . The path, the image of  $T$  under  $P$  is denoted as  $P(T)$ .

A  $T_p$ -tree and the sequence of two ept's reducing it to a path are illustrated in Figure 1.2.




Figure 1.2

**Theorem 2.1.** If  $T$  be a  $T_p$ -tree on  $m$  vertices, then the graph  $T \hat{O} nC_4$  is a vertex equitable graph.

**Proof:** Let  $T$  be a  $T_p$ -tree with  $m$  vertices. By the definition of a transformed tree there exists a parallel transformation  $P$  of  $T$  such that for the path  $P(T)$  we have (i)  $V(P(T)) = V(T)$  (ii)  $E(P(T)) = (E(T) - E_d) \cup E_p$  where  $E_d$  is the set of edges deleted from  $T$  and  $E_p$  is the set of edges newly added through the sequence  $P = (P_1, P_2, \dots, P_k)$  of the epts  $P$  used to arrive at the path  $P(T)$ . Clearly,  $E_d$  and  $E_p$  have the same number of edges.

Now denote the vertices of  $P(T)$  successively by  $u'_1, u'_2, \dots, u'_m$  starting from one pendant vertex of  $P(T)$  right up to the other one. Let  $u_{i1}, u_{i2}, \dots, u_{i(n+1)}, v_{i1}, v_{i2}, \dots, v_{in}$  and  $w_{i1}, w_{i2}, \dots, w_{in}$  ( $1 \leq i \leq m$ ) be the vertices of  $i^{\text{th}}$  copy of  $P_n$  with  $u_{i(n+1)} = u'_i$ . Then  $V(T \hat{O} nC_4) = \{u_{ij} : 1 \leq i \leq m, 1 \leq j \leq n+1 \text{ with } u_{i(n+1)} = u'_i\} \cup \{u'_i, v_{ij}, w_{ij} : 1 \leq i \leq m, 1 \leq j \leq n\}$  and  $E(T \hat{O} nC_4) = E(T) \cup E(nC_4)$ . Here  $|V(T \hat{O} nC_4)| = m(3n+1)$  and  $|E(T \hat{O} nC_4)| = 4mn + m - 1$ . Let  $A = \{0, 1, 2, \dots, \left\lceil \frac{4mn + m - 1}{2} \right\rceil\}$ . Define a vertex labeling  $f: V(T \hat{O} nC_4) \rightarrow A$  as follows.

$$\text{For } 1 \leq i \leq m, 1 \leq j \leq n+1 \quad f(u_{ij}) = \begin{cases} \frac{(4n+1)(i-1)}{2} + 2(j-1) & \text{if } i \text{ is odd} \\ \frac{(4n+1)i}{2} - 2(j-1) & \text{if } i \text{ is even} \end{cases}.$$

For  $1 \leq i \leq m, 1 \leq j \leq n$

$$f(v_{ij}) = \begin{cases} \frac{(4n+1)(i-1)}{2} + 2j & \text{if } i \text{ is odd} \\ \frac{(4n+1)i}{2} - 2j & \text{if } i \text{ is even} \end{cases}, \quad f(w_{ij}) = \begin{cases} \frac{(4n+1)(i-1)}{2} + 2j - 1 & \text{if } i \text{ is odd} \\ \frac{(4n+1)i}{2} - (2j-1) & \text{if } i \text{ is even} \end{cases}.$$

Let  $v_i v_j$  be a transformed edge in  $T$  for some indices  $i, j, 1 \leq i \leq j \leq m$ . Let  $P_1$  be the ept that deletes the edge  $v_i v_j$  and adds an edge  $v_{i+t} v_{j-t}$  where  $t$  is the distance of  $v_i$  from  $v_{i+t}$  and the distance of  $v_j$  from  $v_{j-t}$ . Let  $P$  be a parallel transformation of  $T$  that contains  $P_1$  as one of the constituent epts.

Since  $v_{i+t} v_{j-t}$  is an edge in the path  $P(T)$ , it follows that  $i + t + 1 = j - t$  which implies  $j = i + 2t + 1$ . Therefore,  $i$  and  $j$  are of opposite parity, that is,  $i$  is odd and  $j$  is even or vice-versa.

The induced label of the edge  $v_i v_j$  is given by  $f^*(v_i v_j) = f^*(v_i v_{i+2t+1}) = f(v_i) + f(v_{i+2t+1}) = (4n+1)(i+t)$

and  $f^*(v_{i+t} v_{j-t}) = f^*(v_{i+t} v_{i+t+1}) = f(v_{i+t}) + f(v_{i+t+1}) = (4n+1)(i+t)$

Therefore,  $f^*(v_i v_j) = f^*(v_{i+t} v_{j-t})$ .

It can be verified that the induced edge labels of  $T \hat{\circ} nC_4$  are  $1, 2, 3, \dots, 8mn+2m-2$  and  $|v_f(a) - v_f(b)| \leq 1$  for all  $a, b \in A$ . Hence,  $T \hat{\circ} nC_4$  is a vertex equitable graph.

An example for the vertex equitable labeling of  $T \hat{\circ} 2C_4$  where  $T$  is a  $T_p$ -tree on 8 vertices is shown in Figure 2.2.




Figure 2.2.

**Theorem 2.2.** If  $G$  is a zig-zag triangle, then  $G$  is a vertex equitable graph.

**Proof:** Let zig-zag triangles be defined as in definition 1.3. Here  $|V(G)| = 2n$  and  $|E(G)| = 4n-3$ . Let  $A = \{0, 1, 2, \dots, \left\lceil \frac{4n-3}{2} \right\rceil\}$ . Define a vertex labeling  $f: V(G) \rightarrow A$  as follows.  $f(y_1)=1, f(x_1)=0$ ,

$$f(x_{4i-2}) = 8i - 5, f(y_{4i-2}) = 8i - 6 \text{ if } 1 \leq i \leq \left\lfloor \frac{n+2}{4} \right\rfloor$$


$$f(x_{4i-1}) = 8i - 4, f(y_{4i-1}) = 8i - 3 \text{ if } 1 \leq i \leq \left\lfloor \frac{n+1}{4} \right\rfloor$$

$$f(x_{4i}) = 8i - 2, f(y_{4i}) = 8i \text{ if } 1 \leq i \leq \left\lfloor \frac{n}{4} \right\rfloor$$

For  $1 \leq i \leq \left\lfloor \frac{n-1}{4} \right\rfloor$  and  $n > 4$   $f(x_{4i+1}) = 8i + 1, f(y_{4i+1}) = 8i - 1$ .

It can be verified that the induced edge labels of  $G$  are  $1, 2, \dots, 4n-3$  and  $|v_f(a) - v_f(b)| \leq 1$  for all  $a, b \in A$ . Hence a zig-zag triangle is a vertex equitable graph.

An example for the vertex equitable labeling of zig-zag triangle is shown in Figure 2.3.



**Figure 2.3**

## REFERENCES

- [1]. F.Harary, Graph theory, Addison Wesley, Massachusetts, 1972.
- [2]. Joseph A.Gallian, A Dynamic Survey of graph labeling, *The Electronic Journal of Combinatorics*, DS6, 2016.



- [3]. A.Lourdusamy and M.Seenivasan, Vertex equitable labeling of graphs, *Journal of Discrete Mathematical Sciences & Cryptography*, Vol.11, No.6 (2008), .727-735.
- [4]. P.Jeyanthi and A.Maheswari, Some Results on Vertex Equitable Labeling, *Open Journal of Discrete Mathematics*, 2(2012), 51-57.
- [5].P.Jeyanthi and A.Maheswari, Vertex equitable labeling of cycle and path related graphs, *Utilitas Mathematica*, 98(2015), 215-226.
- [6].P.Jeyanthi and A.Maheswari, Vertex equitable labeling of Transformed Trees, *Journal of Algorithms and Computation* 44 (2013), 9 - 20.
- [7].P.Jeyanthi and A.Maheswari, Vertex equitable labeling of cyclic snakes and bistar graphs, *Journal of Scientific Research*, Vol. 6, No.1(2014), 79-85.
- [8] .P.Jeyanthi ,A.Maheswari and M.Vijayalakshmi *Vertex equitable labeling of cycle and star related graphs*, *Journal of Scientific Research*, Vol. 7, No3(2015), 33-42.
- [9]..P.Jeyanthi A.Maheswari and M.Vijaya Lakshmi, Vertex equitable labeling of double alternate snake graphs *Journal of Algorithms and Computation*, 46 (2015) PP. 27 – 34.
- [10].P.Jeyanthi A.Maheswari, and M.Vijaya Lakshmi , New results on vertex equitable labeling, *Journal of Algebra Combinatorics Discrete structures and Applications* Vol 3(2) (2016) 97-104..
- [11].P.Jeyanthi A.Maheswari, and M.Vijaya Lakshmi , Vertex Equitable Labeling of Super Subdivision Graphs *Scientific International*, 27(4)(2015), 1-3.
- [12].P.Jeyanthi, A.Maheswari and M.Vijaya Lakshmi, Vertex Equitable Labeling of Union of Cyclic Snake graphs, *Proyecciones Journal of Mathematics*, Vol. 35, No2, pp. 177-186, June 2016.
- [13].C.Barrientos, Graceful labeling of cyclic snakes , *Ars Combinatoria* , 60(2001) , 85-96.
- [14]. S. M. Hegde, and Sudhakar Shetty, On Graceful Trees, *Applied Mathematics E- Notes*, 2, pp. 192-197, (2002).